

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to average 2 hours per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, sexual orientation, or marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.

**U.S. DEPARTMENT OF AGRICULTURE
AGRICULTURAL MARKETING SERVICE
SCIENCE AND TECHNOLOGY
PLANT VARIETY PROTECTION OFFICE
BELTSVILLE, MD 20705**

**OBJECTIVE DESCRIPTION OF VARIETY
BUFFALOGRASS (*Buchloe dactyloides*)**

| | | |
|---|---------------------------------------|-----------------------|
| NAME OF APPLICANT (S) | TEMPORARY OR EXPERIMENTAL DESIGNATION | VARIETY NAME |
| ADDRESS (Street and No. or RD No., City, State, Zip Code and Country) | | FOR OFFICIAL USE ONLY |
| | | PVPO NUMBER |

PLEASE READ ALL INSTRUCTIONS CAREFULLY:

Place the appropriate number that describes the varietal character of this variety in the boxes below. Fill unused columns with zeroes (e.g., when the number is 99). In comparisons to the standard varieties, the value should only be used to indicate that the varieties are equal. Characteristics described, including numerical measurements, should represent those that are TYPICAL for the variety. Measured data should be for SPACED PLANTS. Characters in item 3 are considered to reflect homogeneity; frequencies of nontypical plants should be taken into regard in Exhibit A. Any recognized color fan, e.g., Natural Bureau of Standards Circular 553 Supplement, may also be used to determine plant colors; designate system used: _____ . Ranges of values may be included with additional description elsewhere in the application.

Note: For single plant data a minimum of 100 plants is suggested.

1. PLOIDY:

1 = Diploid (2N=14) 2 = Tetraploid (2N=28) 3 = Other _____

2. ADAPTATION: (for forage or pasture)

1 = Northeast 2 = East Central 3 = Southeast 4 = North Central
5 = South Central 6 = Pacific NW. 7 = Southwest 8 = Other _____

3. WINTER HARDINESS:

3 = Tender 5 = Intermediate 7 = Hardy

4. PLANT HEIGHT: (from soil to level of top of panicle)

cm Tall

Compared to Texoka cm (Shorter) (Taller)

5. PLANT GROWTH TYPE: (at maturity)

Type: 1 = Prostrate 2 = Intermediate (Texoka) 3 = Erect (Comanche)

Stolon growth – plants 2 years old compared to comparable plants

cm Stolon Length Compared to Texoka cm (Shorter) (Taller)

Inflorescence Structural Differences

1 = Dioecious 2 = Monoecious 3 = Hermaphroditic

6. LEAF:

Leaf Color: 1 = Light Green 2 = Green (Texoka) 3 = Dark Green 4 = Blue Green (Comanche)

Leaf Hairiness (% Plants with each surface)

% Glabrous % Slightly Pubescent % Pubescent

mm Width (First leaf blade); Compared to Texoka mm (Narrower) (Wider)

mm Length (First leaf blade); Compared to Texoka mm (Shorter) (Longer)

7. BURRS: (seed)

mm Width; Compared to Texoka mm (Narrower) (Wider)

mm Length; Compared to Texoka mm (Shorter) (Longer)

mg per 1,000 pure seed; Compared to Texoka mg (Lighter) (Heavier)

8. DISEASE AND INSECT RESISTANCE: (rate resistance 0-9, where 0 = not tested, 1 = 100% susceptible, and 9 = 100% resistant):

- Powdery Mildew (*Erysiphe Graminis*)
- Common Root Rot (*Rhizotonia Solani Pyhium spp.*)
- Anthracnose (*Colletotrichum Graminicola*) Other _____
- Ergot (*Claviceps Purpurea*)
- Buffalograss Mite (*Eriophyes Aceria*)

Rust and leaf spot: Specify as completely as possible including species and races where known. If generalized resistance or susceptibility is claimed (first box), include or append explanation. (0 = Not Tested, 1 -9 = 100% Susceptibility to 100% Resistant, Respectively).

Comments:

- Leaf Spot.....
- Purple Leaf Spot (*Stagonospora Maculata*).....
- Scald (*Phynchosporium Orthosporium*).....
- Leaf Spot (*Mastigosporium Rubicosum*).....
- Leaf Spot (*Helminthosporium spp.*)
- Other

10. Indicate the Variety that most closely resembles the application variety for the following characters:

| CHARACTER | VARIETY | CHARACTER | VARIETY |
|------------------|---------|----------------|---------|
| Leafiness | | Seedling Vigor | |
| Winter Hardiness | | Seed Size | |
| Stolon Growth | | Persistence | |
| Summer Dormancy | | Leaf Color | |
| Heat Tolerance | | | |

References: Comparative turfgrass evapotranspiration rates and associated plant morphology characteristics K.S. Kim and J. B. Beard Crop. Sci. 28:328-331 (1988). Sex expression in buffalograss under different environments. D. R. Huff and Lin Wu Crop Sci. 27:623-626 (1987)

COMMENTS: